

Gröbner Basis Convex Polytopes and Planar Graph

Dang Vu Giang

Hanoi Institute of Mathematics

18 Hoang Quoc Viet, 10307 Hanoi, Vietnam

e-mail: <dangvugiang@yahoo.com>

February 22, 2012

Abstract. Using the Gröbner basis of an ideal generated by a family of polynomials we prove that every planar graph is 4-colorable. Here we also use the fact that the complete graph of 5 vertices is not included in any planar graph.

Keywords: complete graph of 5 vertices, generators of an ideal, polynomial ring

AMS subject classification: 13P10

Consider a planar graph G with n vertices. Between two any vertices there is at most one edge. We want to color the vertices of G by 4 colors such that no two vertices connected by an edge are colored the same way. We let $\zeta = i \in \mathbb{C}$ be the 4th primitive root of unity ($i^4 = 1$). Let $x_1, x_2, \dots, x_n$ be variables representing the distinct vertices of the graph G . Each vertex is to be assigned one of colors $1, i, i^2, i^3$. This means that $x_k^4 - 1 = 0$ for $k = 1, 2, \dots, n$. Moreover, if x_j and x_k are connected then $x_j^3 + x_j^2 x_k + x_j x_k^2 + x_k^3 = 0$. Let I be the ideal generated by $P_k = x_k^4 - 1$ for $k = 1, 2, \dots, n$ and $Q_{j,k} = x_j^3 + x_j^2 x_k + x_j x_k^2 + x_k^3$ if x_j and x_k are connected. We should prove that $I \neq \mathbb{C}[x_1, x_2, \dots, x_n]$. In fact, the complete graph of 5 vertices

is not included in any planar graph. Not every $Q_{j,k}$ is a generator of I . Therefore, the set of generators of I is not so large. Now we induct on n . Assume that every planar graph of $n - 1$ vertices is 4-colorable. Now consider a planar graph G of n vertices. If some vertex of G has degree less than 4 then G is clearly, 4-colorable. Hence, we assume that vertex of G has degree larger than 3. Let x_n be the vertex of smallest degree and let $x_1, x_2, \dots, x_k$ be vertices connected with x_n . Here, k is the degree of the vertex x_n . Thus, $k \geq 4$. Without loss of generality we assume that $x_1, x_2, \dots, x_k$ are colored by 4 distinct colors. Now assume for the sake of a contradiction that $I_n = \mathbb{C}[x_1, x_2, \dots, x_n]$. That is a real ideal I_{n-1} of $\mathbb{C}[x_1, x_2, \dots, x_{n-1}]$ together with $P_n = x_n^4 - 1$ and $Q_{j,n} = x_n^3 + x_n^2 x_j + x_n x_j^2 + x_j^3$ for $j = 1, 2, \dots, k$ generates the whole ring $\mathbb{C}[x_1, x_2, \dots, x_n]$. We can write $1 = (x_n^4 - 1) \varphi_0 + \sum_{j=1}^k (x_n^3 + x_n^2 x_j + x_n x_j^2 + x_j^3) \varphi_j + f(x_1, x_2, \dots, x_{n-1})$ where $f \in I_{n-1}$ and $\varphi_j \in \mathbb{C}[x_1, x_2, \dots, x_n]$ for $j = 0, 1, 2, \dots, k$.

Acknowledgement. Deepest appreciation is extended towards the NAFOSTED (the National Foundation for Science and Technology Development in Vietnam) for the financial support.

References

- [1] W. Adams and P. Loustaunau, An introduction to Gröbner basis, AMS 1996.
- [2]